Unlocking the Neurogenic Potential of Mammalian Müller Glia

Xiaohuan Xia, Iqbal Ahmad

Department of Ophthalmology and Visual Sciences, University of Nebraska Medical Center, Omaha, NE, USA

Müller glia (MG) are the primary support cells in the vertebrate retina, regulating homeostasis in one of the most metabolically active tissues. In lower vertebrates such as fish, they respond to injury by proliferating and reprogramming to regenerate retinal neurons. In mammals, MG may also react to injury by proliferating, but they fail to initiate regeneration. The barriers to regeneration could be intrinsic to mammalian MG or the function of the niche that cannot support the MG reprogramming required for lineage conversion or both. Understanding these mechanisms in light of those being discovered in fish may lead to the formulation of strategies to unlock the neurogenic potential of MG and restore regeneration in the mammalian retina.

Keywords: Müller glia, Regeneration, Neurogenesis, miR-124, Lin28, Retina

Introduction

Recent progress in our understanding of brain development has significantly altered concepts for treating neurodegenerative diseases, including those that affect the retina to cause blindness. Contrary to previous thought, it is now recognized that neurons are generated in the sub ventricular zone (SVZ) of the lateral ventricle, and the sub-granular zone (SGZ) in the hippocampus from neural stem cells (NSC) of glial origin throughout life (1). Outside these discrete regions in the mammalian CNS, including the retina, active neurogenesis has not been reproducibly demonstrated under normal conditions (2). However, rare neurogenic changes are observed in the injured adult mammalian retina; the source of in-

jury-induced neurogenesis is traced to Müller glia (MG) (3-5). Recent observations that mammalian MG possess NSC properties and are able to generate retinal neurons *in vitro* and upon transplantation *in vivo* posit these cells as the potential target for regenerating diseased or injured photoreceptors and retinal ganglion cells (6). While growing evidence confirms that MG possess evolutionarily conserved neurogenic potential, studies from various labs have observed that, unlike their lower vertebrate counterparts, the injury or disease-activated mammalian MG proliferate, but that they convert to neurons rather infrequently (4, 7, 8). Whether or not the converted neurons are functional, make synaptic connections, and survive for the long term remains unknown. The challenge that remains is how to unlock the neurogenic potentials of the mammalian MG *in vivo*. This may essentially involve approaches to reprogram these cells to dedifferentiate and regain their lost ability to generate neurons or trans-differentiate them into neurons.

Neurogenic potentials of MG in mammals

To address this challenge, the neurogenic potentials of MG need to be evaluated against the backdrop of two models, with the understanding that examples from lower vertebrates would constitute a framework, but that sol-

Accepted for publication July 20, 2016, Published online November 30, 2016
Correspondence to **Iqbal Ahmad**

Department of Ophthalmology and Visual Sciences, University of Nebraska Medical Center, 4046 DRC1, 985840 Omaha, NE 68198-5840, USA

Tel: +402-559-4091, Fax: +402-559-3238

E-mail: iahmad@unmc.edu

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

utions will be unique to the mammalian retina: (1) SVZ/SGZ model, where MG have NSC properties and thus the *capacity* for neuronal differentiation like radial glia-derived NSCs in SVZ and SGZ. Given the observations that the adult NSCs are responsive to environmental cues for neuronal differentiation (1), MG in this model should be amenable to directed differentiation along neuronal lineage through the manipulation of niche-based signals by recombinant growth factors and/or small molecules. (2) Extra SVZ/SGZ model, where MG may have neurogenic *potential* like parenchymal astrocytes but incapable of neuronal differentiation, presumably under the influence of the non-neurogenic niche. For example, both MG and astrocytes express NSC regulators such as Sox2 (MG) and Pax6 (astrocytes) and when cultured in the presence of mitogens tend to differentiate along neuronal lineage (2, 6, 9, 10). However, *in vivo*, although they proliferate and express neural progenitor markers such as Nestin in response to injury, they maintain their glial phenotype, with rare expression of neuronal markers (2, 5, 10). Given their stable glial identity for supporting neurons despite expressing some of the NSC-specific genes, a trans-differentiation approach for the lineage conversion of MG has emerged as a practical option in this

model. This notion is supported by recent observations that injury-activated resident astrocytes differentiate into functional neurons upon enforced expression of NSC regulator, *Sox2* and/or proneural gene, *Ascl1* (11). Regardless, information gained in deciphering the two models will be useful. In both cases, knowledge of the developmental mechanism for the lineage switch, the reaction of MG to different kinds of injuries, phenotypes of the activated MG in spatial and temporal contexts, transcriptional and epigenetic status, and niche-related signals will be critical. These approaches will reveal whether the roadblock to regeneration is cell-intrinsic or cell-extrinsic or both, information that will be essential in formulating strategies for making MG-dependent therapeutic regeneration practical and clinically applicable. Toward this effort, the following issues should be addressed (Fig. 1).

Mechanisms underlying the neuroglial switch in the mammalian retina

The recent success of reprogramming to induce pluripotency in somatic cells and lineage-specific differentiation of pluripotent cells, whether through directed differentiation (SVZ/SGZ model) or trans-differentiation (Extra SVZ/SGZ model), is owed directly to the knowl-

Fig. 1. A schematic representation of mechanisms underlying the neurogenic potential of MG. A subset of MG, most likely those with dormant stem cell properties, respond to injury by proliferating presumably under Notch signaling influence. As the activated MG migrate out of the INL, the changing niche, presumably reflected in altered Notch/cytokine signaling, influence them to engage both the excitatory (Green) and inhibitory (Red) axes regulating the neurogenic potential. It is possible that the imbalance between the two axes and their inadequate niche-based recruitment prevents mammalian MG from regenerating retinal neurons. The niche could be composed of retinal neurons, immigrant astrocytes, microglia, and endothelial cells. The niche-based communication for regeneration may involve diverse signaling pathways, exemplified by Notch and cytokine signaling, acting in concert. Notch signaling besides influencing these two molecular axes, may directly influence the expression of proneural genes. ONL: outer nuclear layer, OPL: outer plexiform layer, INL: inner nuclear layer, IPL: inner plexiform layer, GCL: ganglion cell layer.

edge of developmental mechanisms (12). There is a significant knowledge gap in our understanding of how neurogenesis shifts to the generation of MG in the mammalian retina, information essential to navigate the cellular or molecular roadblocks to neuronal differentiation. For example, although we know that Notch signaling regulates the generation of MG, how it is incorporated into the gliogenic program and to what extent it is involved in suppressing neuronal differentiation remains rather unknown (4). Given the contextual role of Notch signaling (4), its complexity due to the oscillatory expression of its effector *Hes1* (13), and its gliogenic interactions with the eye field gene *Lhx2* (14), this information will be essential to formulate niche-based approaches for facilitating neuronal differentiation of MG. Along this line, it will also be advantageous to know whether genes involved in neurogliogenesis elsewhere in the CNS, such as *Lin28*, a heterochronic gene regulating the developmental timing (15) and *Ezh2*, an epigenetic regulator of gene expression (16) participate in MG differentiation and coordinate the influence of the niche. Recent observations that the decision of retinal progenitors during late histogenesis to differentiate along glial or neuronal lineage is influenced by *Lin28* (17) and that *Ezh2* plays a role in constraining the generation of MG (18) posit these genes as potential targets for pivoting MG towards neuronal differentiation.

Molecular axes involved in MG-mediated regeneration

Regardless of whether or not MG possesses dormant stem cell properties according to the aforementioned models they must initiate and complete an intertwined program of activation and neuronal conversion for successful regeneration (4). Given that injuries in general lead to proliferation of MG across species, but not to their neuronal differentiation in higher vertebrates, it is likely that the cross-talk between transcriptional networks subserving the activation and neuronal differentiation are either not connected to the process, or the network components are in place but are not epigenetically primed for optimal expression in the mammalian MG. These probabilities could be examined by a genome-wide screening of prospectively enriched MG in select animal models in quiescent and activated states (facilitated by lineage reporters and other enrichment protocols such as the side population (SP) cell profiling by microarrays, RNA seq, and ChIP seq analyses. A similar approach to characterizing MG in different states of activation and differentiation in controlled conditions *in vitro* will yield corroborative evidence and facilitate the means to test putative mechanism(s) by manipulating gene expression. Together, these

approaches may provide insight into molecular axes that can be tested against the background of those operational in zebrafish (7, 8, 19). For example, the molecular axis defined by *Lin28* and *Ascl1*, which facilitates MG-mediated regeneration in zebrafish, is a valid target for study in mammals. A variety of approaches in higher vertebrates have demonstrated important regulatory roles for *Lin28* in the maintenance of neural progenitors and neuroglial decision (20). It is likely that *Lin28* influences different developmental function by contextual recruitment of *Hmga2*, a gene encoding a DNA architecture protein (21, 22) and *Ascl1* (23) by directly regulating the heterochronic miRNA *let-7* (17, 21, 23). *let-7* targets *Hmga2* (17, 21) and *Ascl1* (23); therefore, the *Lin28-let-7-Hmga2/Ascl1* axis has emerged as an evolutionary conserved axis that could be a candidate for unlocking neurogenic potential of MG. Early evidence supports this premise. For example, overexpression of *Lin28a* (24) in the enriched MG (Fig. 2A) prompted these cells to acquire neuronal morphology and to express immunoreactivities and transcripts corresponding to neuronal genes (Fig. 2B). More importantly, neuronal differentiation was accompanied by an increase in levels of *miR-124*, a proneural miRNA (25), *Hmga2*, and *Ascl1*, and a decrease in those of *REST*, a global inhibitor of neuronal differentiation (26), thus demonstrating the capacity of *Lin28a* in activating the neurogenic program in MG. However, targeting the *Lin28-let-7-Hmga2/Ascl1* axis alone may not be sufficient for the functional reprogramming of MG along neuronal lineage, given that its influence may not be able to completely counter the inhibitory resistance of the *REST* axis. *REST*, by suppressing the expression of proneural miRNAs, *miR-124* and *miR-9-9**, whose targets are proglial genes encoding SOX9 and HES family of regulators, may keep MG non-neuronal and therefore non-regenerative (25). The existence of the *REST-miR-124-9-9*-Sox9/Hes1* axis in MG suggested that either inhibiting *REST* or ectopically expressing *miR-124/miR-9-9** or both may direct MG along the neuronal lineage. It is likely that the ectopic expression of *miR-124/miR-9-9** in itself might be effective, as they directly or indirectly influence the expression and function of *REST* (25). This notion was supported by the following observations. For example, over expression of *miR-124-9-9** (27) in MG (Fig. 2A) facilitated neuronal morphology with accompanied expression and suppression of neuronal- and glial-specific genes, respectively (Fig. 2C). Expression of both *Ascl1* and *Lin28a* was up regulated and that of *REST* was down regulated in the perturbed groups, compared to controls. However, no significant difference in *Hmga2* transcript levels was observed, suggesting a

Fig. 2. The intrinsic molecular axes and their involvement in the neurogenic potential of MG. (A) A schematic diagram shows the virus-mediated ectopic expression in enriched MG to perturb *Lin28a* (*Lin28a-let7-Hmga2/Ascl1*) and *miR-124* and *miR-9/9** (*Rest-miR-124/miR-9-9*-Sox9/Hes*). (B) *Lin28a* overexpression caused a significant decrease and increase in transcripts characterizing MG (*Glast*) and neurons (*beta-tubulin* and *Map2*), respectively. Neuronal differentiation of transduced MG could also be ascertained by their neuronal morphology and expression of immunoreactivities corresponding to beta-tubulin and Map2. Neuronal differentiation of MG was accompanied by a significant increase and decrease in the expression of positive (*miR-124*, *Hmga2*, and *Ascl1*) and negative (*REST* and *Hes1*) regulators of neurogenesis. (C) Similarly, *miR-124-9-9** overexpression led to a decrease in levels of glia-specific (*Glast*) and an increase in neuron-specific (*beta-tubulin* and *Map2*) transcripts. As observed in *Lin28a* overexpression, transduced MG displayed neuronal morphology and expressed beta-tubulin and Map2 immunoreactivities. A significant increase and decrease in the expression of positive (*Lin28a* and *Ascl1*) and negative (*REST* and *Hes1*) regulators of neurogenesis was accompanied by *miR-124-9-9** mediated neuronal differentiation of MG. Levels of *Hmga2* transcripts remained unchanged. That the two axes might intersect and influence each other was demonstrated by the increase of *miR-124* transcript levels in *Lin28a* overexpression experiment and vice versa (enclosed graph, B, C). MG were enriched as described in Das et al., (6). They were transduced with *Lin28a* retrovirus (24) or *miR124-9-9** lentivirus (27) and the effects of perturbations on gene expression and phenotype were examined by Q-PCR and immunocytochemical analyses, respectively, as described in Parameswaran et al., (22) and Xia et al., (17). Controls included MG transduced with empty retrovirus/lentivirus. Levels of transcripts and miRNA are presented after normalization with that of *GAPDH* and *U6*, respectively. Arrowheads indicate transduced MG (GFP⁺/RFP⁺) expressing neuronal markers, beta-tubulin or Map2. Scale bar, 50 μ m. Data are mean $\pm$ s.e.m.

threshold requirement of *Lin28a* levels for influencing *Hmga2* expression. That *miR-124* and *miR-9-9** could facilitate proneural function of *Ascl1* was demonstrated by the improvement of *Ascl1*-mediated reprogramming of MG by ectopically expressed miRNAs (28). Together, these observations posit these molecular axes as valid intrinsic targets for studying and unlocking the dormant regenerative potential of MG.

Influences of the niche and how they are mediated

Regeneration takes place when homeostasis is disturbed. MG, the guardian of homeostasis in the retina, respond to injuries by proliferating and migrating out of the inner nuclear layer (5). However, despite proliferation and migration as in zebrafish retina, as mentioned, mammalian MG do not initiate regeneration effectively. This raises the possibility that, in addition to internal constraints discussed above, the environment in the mammalian retina might not be conducive for neurogenic conversion of MG. A niche-based approach to unlock neurogenic potential of MG will involve examining the relationship of MG with neighboring retinal cells, microglia, immigrant astrocytes, and endothelial cells in the context of signaling pathways and their capacity to engage the molecular axes involved in reprogramming along neuronal lineage. For example, the understanding of the role of microglia, one of the first responders to injury, and which may mediate regeneration through inflammatory signals, is still evolving in the retina (29). Studies have demonstrated that Notch (8, 30), Wnt (5, 6), FGF (6, 8), insulin and insulin-like growth factor-1 (8), Shh (31), and cytokines such as TNF α (8, 19), leptin, and IL-11 (32) may play important roles in mediating the influence of the niche in reprogramming MG. However, the identity of cells delivering the signals and whether or not these signals and their associated pathways act in concert, which would determine niche-based strategies to promote regeneration, remain largely unknown. However, studies in zebrafish have begun to shed light on these issues. For example, it has been observed that TNF α released by dying photoreceptors in mechanically damaged zebrafish retina may constitute one of the early signals for activating MG for regeneration (19). Also, it was reported that zebrafish MG respond to retinal injury by secreting leptin and IL-11, which help reprogram cells in an autocrine fashion (32). In both cases, these cytokines facilitated injury-dependent induction of *Ascl1*, a key step in reprogramming of MG (8). Whether or not signaling-mediated by these factors are similarly involved in the mammalian retina remains to be demonstrated. Although the identity of cells delivering Notch signaling in MG re-

mains speculative, emerging evidence from zebrafish and higher vertebrates posit it as an important niche-based target for MG-dependent regeneration. It remains active long after facilitating the differentiation of MG, enabling them to launch the proliferative response when injury takes place (4, 5). However, persistence of Notch signaling in activated MG, which are poised for neuronal conversion, might be deleterious for regenerative process, given the inhibitory influence of Notch signaling on the activation of proneural genes (33). The inability to confine the activation of Notch signaling in MG to a narrow temporal window following injury may be one of the reasons for their anemic neuronal conversion in mammals, compared to that in zebrafish (30). Emerging evidence, once again from the zebrafish model, suggests that these pathways, including Notch signaling, may act in concert and MG integrate the diverse niche-based influence of a variety of signaling molecules for a calibrated regenerative response to injury (8). Therefore, niche-related information is crucial for understanding the temporal and spatial modulation of signaling required for activating MG and shifting them from the state of activation to neuronal differentiation.

Markers of activated MG

This is fundamental information missing from studies of MG-dependent regeneration, without which the underlying mechanisms remain difficult to grasp and the therapeutic target rather illusory. Currently, the identity of the activated MG represents cells in the inner nuclear layer of the retina, which have incorporated BrdU in response to injury and express MG-specific markers. Given the transition of MG from a quiescent to a proliferative state and the generation of their progenies in different stages of development, the identity based on BrdU incorporation, though important, is of limited value. Attempts have been made to prospectively enrich activated MG by Hoechst dye efflux assays and to characterize them as SP cells, a phenotype shared by the majority of stem cells (4, 6). However, since SP cells are also heterogeneous, this functional approach has limitations in the absence of specific cellular markers. Although several signaling pathways (e.g. Notch signaling) and intrinsic factors (e.g., *Ascl1*) have been observed to regulate MG activation in fish, birds, and rodents (8), their status as markers of activated MG in different stages of regeneration remain unspecified. Progress in understanding the regenerative mechanisms in tissues like blood (34), intestine (35), and skin (36) has been facilitated by the identification and characterization of markers of the resident stem cells and their progenies. Within the CNS, insight into regeneration

in the SVZ has come from the characterization of B1 quiescent and active progenitors and resulting neuroblasts (37). Therefore, to follow MG-mediated regeneration, spatially and temporally, and to identify the stage-specific intrinsic players, a concerted effort is needed to search for reliable and reproducible markers.

Animal models for MG-mediated therapeutic regeneration

Currently, there is no consensus on reliable and reproducible mammalian models to study the activation and neurogenic potential of MG. Rodent models representing retinal injuries, ranging from those caused by the exposure to light, neurotoxins, and genetic mutations have been used. Given the observations that glia respond differently to different types and durations of injuries (2), that their levels of activation differ from species to species (Ahmad et al., unpublished observations), and that within the same species there are strain differences in responses (38), a case may be made for developing injury- and species/strain-specific models for reproducible and unambiguous examination of MG neurogenic potentials. These models will be valuable in incorporating transgenic technology for reliable lineage tracing, and more importantly for molecular characterization of activated MG. Characterization of these cells for gene regulatory network and epigenetic signature is essential to shed light on the status of their neurogenic potential and approaches to unlock it. Furthermore, cell type specific injury models would test the ability of MG to replace specific neuronal types, which will be helpful in designing approaches for MG-dependent therapeutic regeneration. For example, the loss of vision in two of the intractable blinding diseases, age-related macular degeneration (AMD) and glaucoma, is due to selective loss of photoreceptors and retinal ganglion cells (RGCs), respectively. The neurotoxin injury models where exposure of the retina to N-Methyl-N-nitrosourea (MNU) ablates photoreceptors (39) and NMDA causes the degeneration of RGCs (40) would reveal whether or not MG could differentiate along specific neuronal types. Also, it may shed light on if MG-dependent therapeutic regeneration would be a practical approach to address degenerative changes in AMD and/or glaucoma.

Conclusions

The discipline of MG-based regeneration is a recent one, particularly in mammals. With information on the nature of the activated MG and the molecular axes mediating their response to the niche, aided by reproducible animal models and lineage reporters, a pharmacological re-

cruitment of these endogenous progenitors for therapeutic regeneration is a near possibility. In addition, information emerging from these studies will help us understand how the neurogenic potential in the adult mammalian brain is constrained relative to that of lower vertebrates, currently a significant knowledge gap.

Acknowledgments

This work was supported by Pearson Foundation and Nebraska Department of Health and Human Services (LB606). We are thankful to Dr. Andrew S. Yoo and Dr. George Q. Daley for the pLemir-9-124 and pMSCV-mLin28a plasmids, respectively.

References

1. Gage FH, Temple S. Neural stem cells: generating and regenerating the brain. *Neuron* 2013;80:588-601
2. Dimou L, Götz M. Glial cells as progenitors and stem cells: new roles in the healthy and diseased brain. *Physiol Rev* 2014;94:709-737
3. Karl MO, Reh TA. Regenerative medicine for retinal diseases: activating endogenous repair mechanisms. *Trends Mol Med* 2010;16:193-202
4. Ahmad I, Del Debbio CB, Das AV, Parameswaran S. Müller glia: a promising target for therapeutic regeneration. *Invest Ophthalmol Vis Sci* 2011;52:5758-5764
5. Del Debbio CB, Balasubramanian S, Parameswaran S, Chaudhuri A, Qiu F, Ahmad I. Notch and Wnt signaling mediated rod photoreceptor regeneration by Müller cells in adult mammalian retina. *PLoS One* 2010;5:e12425
6. Das AV, Mallya KB, Zhao X, Ahmad F, Bhattacharya S, Thoreson WB, Hegde GV, Ahmad I. Neural stem cell properties of Müller glia in the mammalian retina: regulation by Notch and Wnt signaling. *Dev Biol* 2006;299:283-302
7. Lenkowski JR, Raymond PA. Müller glia: Stem cells for generation and regeneration of retinal neurons in teleost fish. *Prog Retin Eye Res* 2014;40:94-123
8. Goldman D. Müller glial cell reprogramming and retina regeneration. *Nat Rev Neurosci* 2014;15:431-442
9. Roesch K, Jadhav AP, Trimarchi JM, Stadler MB, Roska B, Sun BB, Cepko CL. The transcriptome of retinal Müller glial cells. *J Comp Neurol* 2008;509:225-238
10. Sakurai K, Osumi N. The neurogenesis-controlling factor, Pax6, inhibits proliferation and promotes maturation in murine astrocytes. *J Neurosci* 2008;28:4604-4612
11. Heinrich C, Bergami M, Gascón S, Lepier A, Viganò F, Dimou L, Sutor B, Berninger B, Götz M. Sox2-mediated conversion of NG2 glia into induced neurons in the injured adult cerebral cortex. *Stem Cell Reports* 2014;3:1000-1014
12. Amamoto R, Arlotta P. Development-inspired reprogramming of the mammalian central nervous system. *Science* 2014;343:1239882
13. Imayoshi I, Isomura A, Harima Y, Kawaguchi K, Kori H,

- Miyachi H, Fujiwara T, Ishidate F, Kageyama R. Oscillatory control of factors determining multipotency and fate in mouse neural progenitors. *Science* 2013;342:1203-1208
14. de Melo J, Zibetti C, Clark BS, Hwang W, Miranda-Angulo AL, Qian J, Blackshaw S. Lhx2 is an essential factor for retinal gliogenesis and notch signaling. *J Neurosci* 2016; 36:2391-2405
15. Balzer E, Heine C, Jiang Q, Lee VM, Moss EG. LIN28 alters cell fate succession and acts independently of the let-7 microRNA during neurogliogenesis in vitro. *Development* 2010;137:891-900
16. Hirabayashi Y, Suzuki N, Tsuboi M, Endo TA, Toyoda T, Shinga J, Koseki H, Vidal M, Gotoh Y. Polycomb limits the neurogenic competence of neural precursor cells to promote astrogenic fate transition. *Neuron* 2009;63:600-613
17. Xia X, Ahmad I. let-7 microRNA regulates neurogliogenesis in the mammalian retina through Hmga2. *Dev Biol* 2016;410:70-85
18. Zhang J, Taylor RJ, La Torre A, Wilken MS, Cox KE, Reh TA, Vetter ML. Ezh2 maintains retinal progenitor proliferation, transcriptional integrity, and the timing of late differentiation. *Dev Biol* 2015;403:128-138
19. Gorsuch RA, Hyde DR. Regulation of Müller glial dependent neuronal regeneration in the damaged adult zebrafish retina. *Exp Eye Res* 2014;123:131-140
20. Rehfeld F, Rohde AM, Nguyen DT, Wulczyn FG. Lin28 and let-7: ancient milestones on the road from pluripotency to neurogenesis. *Cell Tissue Res* 2015;359:145-160
21. Nishino J, Kim I, Chada K, Morrison SJ. Hmga2 promotes neural stem cell self-renewal in young but not old mice by reducing p16Ink4a and p19Arf Expression. *Cell* 2008;135: 227-239
22. Parameswaran S, Xia X, Hegde G, Ahmad I. Hmga2 regulates self-renewal of retinal progenitors. *Development* 2014;141:4087-4097
23. Cimadamore F, Amador-Arjona A, Chen C, Huang CT, Tersikh AV. SOX2-LIN28/let-7 pathway regulates proliferation and neurogenesis in neural precursors. *Proc Natl Acad Sci U S A* 2013;110:E3017-E3026.
24. Viswanathan SR, Powers JT, Einhorn W, Hoshida Y, Ng TL, Toffanin S, O'Sullivan M, Lu J, Phillips LA, Lockhart VL, Shah SP, Tanwar PS, Mermel CH, Beroukhi R, Azam M, Teixeira J, Meyerson M, Hughes TP, Llovet JM, Radich J, Mullighan CG, Golub TR, Sorensen PH, Daley GQ. Lin28 promotes transformation and is associated with advanced human malignancies. *Nat Genet* 2009;41:843-848
25. Stappert L, Roese-Koerner B, Brüstle O. The role of microRNAs in human neural stem cells, neuronal differentiation and subtype specification. *Cell Tissue Res* 2015; 359:47-64.
26. Qureshi IA, Gokhan S, Mehler MF. REST and CoREST are transcriptional and epigenetic regulators of seminal neural fate decisions. *Cell Cycle* 2010;9:4477-4486
27. Yoo AS, Staahl BT, Chen L, Crabtree GR. MicroRNA-mediated switching of chromatin-remodelling complexes in neural development. *Nature* 2009;460:642-646
28. Wohl SG, Reh TA. miR-124-9-9* potentiates Ascl1-induced reprogramming of cultured Müller glia. *Glia* 2016;64:743-762
29. Fischer AJ, Zelinka C, Gallina D, Scott MA, Todd L. Reactive microglia and macrophage facilitate the formation of Müller glia-derived retinal progenitors. *Glia* 2014; 62:1608-1628
30. Del Debbio CB, Mir Q, Parameswaran S, Mathews S, Xia X, Zheng L, Neville AJ, Ahmad I. Notch signaling activates stem cell properties of müller glia through transcriptional regulation and Skp2-mediated degradation of p27Kip1. *PLoS One* 2016;11:e0152025
31. Wan J, Zheng H, Xiao HL, She ZJ, Zhou GM. Sonic hedgehog promotes stem-cell potential of Müller glia in the mammalian retina. *Biochem Biophys Res Commun* 2007; 363:347-354
32. Zhao XF, Wan J, Powell C, Ramachandran R, Myers MG Jr, Goldman D. Leptin and IL-6 family cytokines synergize to stimulate Müller glia reprogramming and retina regeneration. *Cell Rep* 2014;9:272-284
33. Kageyama R, Ohtsuka T, Hatakeyama J, Ohsawa R. Roles of bHLH genes in neural stem cell differentiation. *Exp Cell Res* 2005;306:343-348
34. Mayle A, Luo M, Jeong M, Goodell MA. Flow cytometry analysis of murine hematopoietic stem cells. *Cytometry A* 2013;83:27-37
35. Gracz AD, Magness ST. Defining hierarchies of stemness in the intestine: evidence from biomarkers and regulatory pathways. *Am J Physiol Gastrointest Liver Physiol* 2014; 307:G260-G273
36. Kretzschmar K, Watt FM. Markers of epidermal stem cell subpopulations in adult mammalian skin. *Cold Spring Harb Perspect Med* 2014;4
37. Lim DA, Alvarez-Buylla A. Adult neural stem cells stake their ground. *Trends Neurosci* 2014;37:563-571
38. Suga A, Sadamoto K, Fujii M, Mandai M, Takahashi M. Proliferation potential of Müller glia after retinal damage varies between mouse strains. *PLoS One* 2014;9:e94556
39. Tsubura A, Yoshizawa K, Kuwata M, Uehara N. Animal models for retinitis pigmentosa induced by MNU; disease progression, mechanisms and therapeutic trials. *Histol Histopathol* 2010;25:933-944
40. Li Y, Schlamp CL, Nickells RW. Experimental induction of retinal ganglion cell death in adult mice. *Invest Ophthalmol Vis Sci* 1999;40:1004-1008